

Please check the examination details below before entering your candidate information

Candidate surname

Other names

**Pearson Edexcel
Level 3 GCE**

Centre Number

--	--	--	--	--

Candidate Number

--	--	--	--

Time 1 hour 30 minutes

Paper
reference

9FM0/3A

Further Mathematics

Advanced

PAPER 3A: Further Pure Mathematics 1

You must have:

Mathematical Formulae and Statistical Tables (Green), calculator

Total Marks

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for algebraic manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
– *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 8 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- Good luck with your examination.

Turn over ►

P66798A

©2021 Pearson Education Ltd.

1/1/1/



Pearson

2. (i) Use the substitution $t = \tan \frac{x}{2}$ to prove the identity

$$\frac{\sin x - \cos x + 1}{\sin x + \cos x - 1} \equiv \sec x + \tan x \quad x \neq \frac{n\pi}{2} \quad n \in \mathbb{Z} \quad (5)$$

- (ii) Use the substitution $t = \tan \frac{\theta}{2}$ to determine the exact value of

$$\int_0^{\frac{\pi}{2}} \frac{5}{4 + 2 \cos \theta} d\theta$$

giving your answer in simplest form.

(5)



4.




Figure 2

A small aircraft is landing in a field.

In a model for the landing the aircraft travels in different straight lines before and after it lands, as shown in Figure 2.

The vector \mathbf{v}_A is in the direction of travel of the aircraft as it approaches the field.

The vector \mathbf{v}_L is in the direction of travel of the aircraft after it lands.

With respect to a fixed origin, the field is modelled as the plane with equation

$$x - 2y + 25z = 0$$

and

$$\mathbf{v}_A = \begin{pmatrix} 3 \\ -2 \\ -1 \end{pmatrix}$$

(a) Write down a vector \mathbf{n} that is a normal vector to the field.

(1)

(b) Show that $\mathbf{n} \times \mathbf{v}_A = \lambda \begin{pmatrix} 13 \\ 19 \\ 1 \end{pmatrix}$, where λ is a constant to be determined.

(2)

When the aircraft lands it remains in contact with the field and travels in the direction \mathbf{v}_L .

The vector \mathbf{v}_L is in the same plane as both \mathbf{v}_A and \mathbf{n} as shown in Figure 2.

(c) Determine a vector which has the same direction as \mathbf{v}_L .

(3)

(d) State a limitation of the model.

(1)



5. The parabola C has equation

$$y^2 = 32x$$

and the hyperbola H has equation

$$\frac{x^2}{36} - \frac{y^2}{9} = 1$$

(a) Write down the equations of the asymptotes of H . (1)

The line l_1 is normal to C and parallel to the asymptote of C with positive gradient.

The line l_2 is normal to C and parallel to the asymptote of C with negative gradient.

(b) Determine (4)

(i) an equation for l_1

(ii) an equation for l_2

The lines l_1 and l_2 meet H at the points P and Q respectively.

(c) Find the area of the triangle OPQ , where O is the origin. (4)



6. $\left[\begin{array}{l} \text{The Taylor series expansion of } f(x) \text{ about } x = a \text{ is given by} \\ f(x) = f(a) + (x - a)f'(a) + \frac{(x - a)^2}{2!}f''(a) + \dots + \frac{(x - a)^r}{r!}f^{(r)}(a) + \dots \end{array} \right]$

Given that

$$y = (1 + \ln x)^2 \quad x > 0$$

(a) show that $\frac{d^2y}{dx^2} = -\frac{2 \ln x}{x^2}$ (4)

(b) Hence find $\frac{d^3y}{dx^3}$ (2)

(c) Determine the Taylor series expansion about $x = 1$ of

$$(1 + \ln x)^2$$

in ascending powers of $(x - 1)$, up to and including the term in $(x - 1)^3$

Give each coefficient in simplest form. (3)

(d) Use this series expansion to evaluate

$$\lim_{x \rightarrow 1} \frac{2x - 1 - (1 + \ln x)^2}{(x - 1)^3}$$

explaining your reasoning clearly. (3)



