



KING EDWARD VI  
HANDSWORTH GRAMMAR  
SCHOOL FOR BOYS



KING EDWARD VI  
ACADEMY TRUST  
BIRMINGHAM

# Year 13

## Pure Mathematics

### P2 6 Trigonometric Functions

HGS Maths



Dr Frost Course



Name: \_\_\_\_\_

Class: \_\_\_\_\_

## Contents

[6.1\) Secant, cosecant and cotangent](#)

[6.2\) Graphs of  \$\sec x\$ ,  \$\operatorname{cosec} x\$  and  \$\cot x\$](#)

[6.3\) Using  \$\sec x\$ ,  \$\operatorname{cosec} x\$  and  \$\cot x\$](#)

[6.4\) Trigonometric identities](#)

[6.5\) Inverse trigonometric functions](#)

**Extract from Formulae booklet**

**Past Paper Practice**

**Summary**

## Prior knowledge check

### Prior knowledge check

- 1** Sketch the graph of  $y = \sin x$  for  $-180^\circ \leq x \leq 180^\circ$ . Use your sketch to solve, for the given interval, the equations:

**a**  $\sin x = 0.8$       **b**  $\sin x = -0.4$


← Year 1, Chapter 10

- 2** Prove that  $\frac{1}{\sin x \cos x} - \frac{1}{\tan x} = \tan x$

← Year 1, Chapter 10

- 3** Find all the solutions in the interval  $0 \leq x \leq 2\pi$  to the equation  $3 \sin^2(2x) = 1$ .

← Section 5.5



Trigonometric functions can be used to model oscillations and resonance in bridges. You will use the functions in this chapter together with differentiation and integration in chapters 9 and 12.

## 6.1) Secant, cosecant and cotangent

## Notes

## Worked Example

Find the exact values of:

$$\operatorname{cosec} \frac{\pi}{2}$$

$$\operatorname{cosec} \frac{\pi}{3}$$

$$\operatorname{cosec} \frac{\pi}{4}$$

$$\operatorname{cosec} 150^\circ$$

$$\operatorname{cosec} (-210^\circ)$$

$$\operatorname{cosec} (0)$$

## Worked Example

Find the exact values of:

$$\sec \frac{\pi}{2}$$

$$\sec \frac{\pi}{3}$$

$$\sec \frac{\pi}{4}$$

$$\sec 150^\circ$$

$$\sec (-210^\circ)$$

$$\sec (0)$$

## Worked Example

Find the exact values of:

$$\cot \frac{\pi}{2}$$

$$\cot \frac{\pi}{3}$$

$$\cot \frac{\pi}{4}$$

$$\cot 150^\circ$$


$$\cot (-210^\circ)$$

$$\cot (0)$$




## 6.2) Graphs of $\sec x$ , $\operatorname{cosec} x$ and $\cot x$


# Graphs



# Graphs



# Graphs



## Notes

## Worked Example

Sketch the graph in the interval  $-2\pi \leq x \leq 2\pi$ :

$$y = \operatorname{cosec} \left( x + \frac{\pi}{4} \right)$$

$$y = \cot \left( x - \frac{\pi}{3} \right)$$

## Worked Example

Sketch the graph in the interval  $-2\pi \leq x \leq 2\pi$ :

$$y = 2\operatorname{cosec} x$$

$$y = 3\cot x$$

## Worked Example

Sketch the graph in the interval  $0 \leq x \leq 2\pi$ :

$$y = 6 + \operatorname{cosec} 4x$$

$$y = \cot 3x - 5$$



## Worked Example

State the range of:

$$y = \operatorname{cosec} x, x \in \mathbb{R}, x \neq n\pi, n \in \mathbb{Z}$$

$$y = \cot x, x \in \mathbb{R}, x \neq n\pi, n \in \mathbb{Z}$$

## Worked Example

Find the range of values of  $k$  for which  $2 + 7 \sec x = k$  has no solutions.

Find the range of values of  $k$  for which  $3 \operatorname{cosec} x - 5 = k$  has no solutions

## Worked Example

Find the maximum and minimum of the graph, stating the smallest positive values of  $\theta$  at which they occur:

$$y = \frac{1}{2 + 3 \sec \theta}$$

### 6.3) Using $\sec x$ , $\operatorname{cosec} x$ and $\cot x$

## Notes

## Worked Example

Simplify:

$$\sin \theta \cos \theta (\operatorname{cosec} \theta - \sec \theta)$$

Simplify:

$$\cos \theta \tan \theta \operatorname{cosec} \theta$$

## Worked Example

Prove that:

$$\frac{\tan \theta \sec \theta}{\sec^2 \theta + \operatorname{cosec}^2 \theta} \equiv \sin^3 \theta$$

## Worked Example

Prove that:

$$\operatorname{cosec} x - \sin x \equiv \cos x \cot x$$



## Worked Example

Prove that:

$$(1 + \sin x)(\sec x - \tan x) \equiv \cos x$$

## Worked Example

Solve in the interval  $0 \leq \theta \leq 2\pi$ :

$$\operatorname{cosec} \theta = 1$$

Solve in the interval  $0 \leq \theta \leq 2\pi$ :

$$\cot \theta = \sqrt{3}$$

### Worked Example

Solve in the interval  $0 \leq \theta \leq 2\pi$ :

$$\operatorname{cosec} \theta = 2$$

Solve in the interval  $0 \leq \theta \leq 2\pi$ :

$$\cot \theta = -3$$

## Worked Example

Solve in the interval  $0 \leq \theta \leq 2\pi$ :

$$\operatorname{cosec} \theta = 0$$

Solve in the interval  $0 \leq \theta \leq 2\pi$ :

$$\cot \theta = 0$$

### Worked Example

Solve in the interval  $-180^\circ \leq \theta \leq 180^\circ$ :

$$\operatorname{cosec} \theta = -\sqrt{2}$$

Solve in the interval  $-180^\circ \leq \theta \leq 180^\circ$ :

$$\cot \theta = -\sqrt{3}$$

## Worked Example

Solve in the interval  $0^\circ \leq \theta \leq 360^\circ$ :

$$\frac{1 - \tan x}{1 - \cot x} = 2$$

## 6.4) Trigonometric identities

From C2 you knew:

$$\sin^2 x + \cos^2 x = 1$$

There are just two new identities you need to know:

Dividing by  $\cos^2 x$ :

$$1 + \tan^2 x = \sec^2 x$$

Dividing by  $\sin^2 x$ :

$$1 + \cot^2 x = \operatorname{cosec}^2 x$$

“Prove that  $1 + \tan^2 x \equiv \sec^2 x$ .”

$$\sin^2 x + \cos^2 x \equiv 1$$

$$\frac{\sin^2 x}{\cos^2 x} + \frac{\cos^2 x}{\cos^2 x} \equiv \frac{1}{\cos^2 x}$$

$$\tan^2 x + 1 \equiv \sec^2 x$$

**Fro Tip:** I used to misremember this as “ $1 + \sec^2 x = \tan^2 x$ ”. Then I imagined the Queen coming back from holiday, saying “One is tanned”, i.e. the 1 goes with the  $\tan^2 x$ .

**Fro Tip:** I remember this one by starting with the above, and slapping ‘co’ on front of each trig function.

**Fro Tip:** This has been asked in an exam before! You must explicitly show each term being divided by  $\cos^2 x$ .

## Notes



## Notes

## Worked Example

Prove that:

$$\operatorname{cosec}^2 \theta - \sin^2 \theta \equiv \cos^2 \theta (1 + \operatorname{cosec}^2 \theta)$$

## Worked Example

593a: Solve a trigonometric equation with  $\sec$ ,  $\operatorname{cosec}$  and  $\cot$  given in the form  $\sec(ax + b) = k$  where  $x$  is in degrees.

Solve  $\operatorname{cosec}(2x + 10) = 1.8$  in the interval  $0^\circ < x < 180^\circ$

Give your solution(s) correct to 2 decimal places where appropriate.

  $x =$    $^\circ$

  $x =$    $^\circ$

## Worked Example


### 593d: Solve quadratic equations in terms of sec, cosec or cot


Solve  $2 \operatorname{cosec}^2 \theta - 3 \operatorname{cosec} \theta - 14 = 0$  in the interval  $0^\circ < \theta < 360^\circ$

Give your solution(s) correct to 1 decimal place where appropriate.

  $\theta =$   °

  $\theta =$   °

  $\theta =$   °

  $\theta =$   °

## Worked Example

Prove that:

$$\sec^4 \theta - \tan^4 \theta = \frac{1 + \sin^2 \theta}{1 - \sin^2 \theta}$$

### Worked Example

Given that  $\tan A = -\frac{3}{4}$  and angle  $A$  is obtuse, find the exact values of  $\sec A$  and  $\sin A$

## Worked Example

Given that  $\cos A = \frac{3}{5}$  and angle  $A$  is reflex, find the exact values of  $\tan A$  and  $\operatorname{cosec} A$

## Worked Example

Given that  $x = \operatorname{cosec} \theta + \cot \theta$ , express in its simplest form:




$$x^2 + \frac{1}{x^2} + 2$$



## 6.5) Inverse trigonometric functions

# Graphs

Fully label:



## Notes

## Worked Example

Work out, in radians, the values of:

a)  $\arcsin\left(\frac{\sqrt{2}}{2}\right)$

b)  $\arccos(1)$

c)  $\arctan(-\sqrt{3})$

## Worked Example

Given that  $y = \arcsin x$ ,  $-1 \leq x \leq 1$  and  $0 \leq y \leq \pi$ ,

- a) Express  $\arccos x$  in terms of  $y$
- b) Hence evaluate  $\arcsin x - \arccos x$

## Worked Example

Prove that for  $0 \leq x \leq 1$ ,  $\arcsin x = \arccos \sqrt{1 - x^2}$  and give a reason why this result is not true for  $-1 \leq x \leq 0$

## Extract from Formulae book

### Trigonometric identities

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B} \quad (A \pm B \neq (k + \frac{1}{2})\pi)$$

$$\sin A + \sin B = 2 \sin \frac{A+B}{2} \cos \frac{A-B}{2}$$

$$\sin A - \sin B = 2 \cos \frac{A+B}{2} \sin \frac{A-B}{2}$$

$$\cos A + \cos B = 2 \cos \frac{A+B}{2} \cos \frac{A-B}{2}$$

$$\cos A - \cos B = -2 \sin \frac{A+B}{2} \sin \frac{A-B}{2}$$

Small angle approximations

$$\sin \theta \approx \theta$$

$$\cos \theta \approx 1 - \frac{\theta^2}{2}$$

$$\tan \theta \approx \theta$$

where  $\theta$  is measured in radians

# Past Paper Questions

## A2 2020 Paper 1

Trig – sec, cosec, cot

12.

In this question you must show all stages of your working.

Solutions relying entirely on calculator technology are not acceptable.

(a) Show that

$$\operatorname{cosec} \theta - \sin \theta \equiv \cos \theta \cot \theta \quad \theta \neq (180n)^\circ \quad n \in \mathbb{Z} \quad (3)$$

(b) Hence, or otherwise, solve for  $0 < x < 180^\circ$

$$\operatorname{cosec} x - \sin x = \cos x \cot(3x - 50^\circ) \quad (5)$$



### Exams

- Formula Booklet
- Past Papers
- Practice Papers
- [past paper Qs by topic](#)

Past paper practice by topic. Both new and old specification can be found via this link on [hgsmaths.com](http://hgsmaths.com)

		(8 marks)	
(p)	Dequies $x = 20^\circ$	B1	3'5'
	$x = 112^\circ$	V1	1'1P
	Also $\cos x = \cos(3x - 20^\circ) \Rightarrow x + 180^\circ = 3x - 20^\circ$	M1	3'1
	$x = 52^\circ$	V1	1'1P
	$\cos x = \cos(3x - 20^\circ) \Rightarrow x = 3x - 20^\circ$	M1	3'1'
	$\Rightarrow \cos x \cos x = \cos x \cos(3x - 20^\circ)$ $\operatorname{cosec} x - \sin x = \cos x \cot(3x - 20^\circ)$		
15 (a)	$\frac{\sin \theta}{\cos \theta} = \cos \theta \times \frac{\sin \theta}{\cos \theta} = \cos \theta \cot \theta$ *	V1*	3'1
	$\operatorname{cosec} \theta - \sin \theta = \frac{\sin \theta}{1} - \sin \theta = \frac{\sin \theta}{1 - \sin^2 \theta}$	M1	3'1
	2 marks for next $\operatorname{cosec} \theta = \frac{\sin \theta}{1}$	B1	1'3
Question	Scheme	Marks	AO*




# Summary of Key Points

## Summary of key points


- $\sec x = \frac{1}{\cos x}$  (undefined for values of  $x$  for which  $\cos x = 0$ )
  - $\operatorname{cosec} x = \frac{1}{\sin x}$  (undefined for values of  $x$  for which  $\sin x = 0$ )
  - $\cot x = \frac{1}{\tan x}$  (undefined for values of  $x$  for which  $\tan x = 0$ )
  - $\cot x = \frac{\cos x}{\sin x}$

2 The graph of  $y = \sec x$ ,  $x \in \mathbb{R}$ , has symmetry in the  $y$ -axis and has period  $360^\circ$  or  $2\pi$  radians. It has vertical asymptotes at all the values of  $x$  for which  $\cos x = 0$ .




- The domain of  $y = \sec x$  is  $x \in \mathbb{R}$ ,  $x \neq 90^\circ, 270^\circ, 450^\circ, \dots$  or any odd multiple of  $90^\circ$ .
- In radians the domain is  $x \in \mathbb{R}$ ,  $x \neq \frac{\pi}{2}, \frac{3\pi}{2}, \frac{5\pi}{2}, \dots$  or any odd multiple of  $\frac{\pi}{2}$ .
- The range of  $y = \sec x$  is  $y \leq -1$  or  $y \geq 1$ .

3 The graph of  $y = \operatorname{cosec} x$ ,  $x \in \mathbb{R}$ , has period  $360^\circ$  or  $2\pi$  radians. It has vertical asymptotes at all the values of  $x$  for which  $\sin x = 0$ .



- The domain of  $y = \operatorname{cosec} x$  is  $x \in \mathbb{R}$ ,  $x \neq 0^\circ, 180^\circ, 360^\circ, \dots$  or any multiple of  $180^\circ$ .
- In radians the domain is  $x \in \mathbb{R}$ ,  $x \neq 0, \pi, 2\pi, \dots$  or any multiple of  $\pi$ .
- The range of  $y = \operatorname{cosec} x$  is  $y \leq -1$  or  $y \geq 1$ .

4 The graph of  $y = \cot x$ ,  $x \in \mathbb{R}$ , has period  $180^\circ$  or  $\pi$  radians. It has vertical asymptotes at all the values of  $x$  for which  $\tan x = 0$ .



- The domain of  $y = \cot x$  is  $x \in \mathbb{R}$ ,  $x \neq 0^\circ, 180^\circ, 360^\circ, \dots$  or any multiple of  $180^\circ$ .
- In radians the domain is  $x \in \mathbb{R}$ ,  $x \neq 0, \pi, 2\pi, \dots$  or any multiple of  $\pi$ .
- The range of  $y = \cot x$  is  $y \in \mathbb{R}$ .


5  $\sec x = k$  and  $\operatorname{cosec} x = k$  have no solutions for  $-1 < k < 1$ .

6 You can use the identity  $\sin^2 x + \cos^2 x \equiv 1$  to prove the following identities:

- $1 + \tan^2 x \equiv \sec^2 x$
- $1 + \cot^2 x \equiv \operatorname{cosec}^2 x$


7 The inverse function of  $\sin x$  is called **arcsin**  $x$ .

- The domain of  $y = \arcsin x$  is  $-1 \leq x \leq 1$
- The range of  $y = \arcsin x$  is  $-\frac{\pi}{2} \leq \arcsin x \leq \frac{\pi}{2}$  or  $-90^\circ \leq \arcsin x \leq 90^\circ$



8 The inverse function of  $\cos x$  is called **arccos**  $x$ .

- The domain of  $y = \arccos x$  is  $-1 \leq x \leq 1$
- The range of  $y = \arccos x$  is  $0 \leq \arccos x \leq \pi$  or  $0^\circ \leq \arccos x \leq 180^\circ$



9 The inverse function of  $\tan x$  is called **arctan**  $x$ .

- The domain of  $y = \arctan x$  is  $x \in \mathbb{R}$
- The range of  $y = \arctan x$  is  $-\frac{\pi}{2} < \arctan x < \frac{\pi}{2}$  or  $-90^\circ < \arctan x < 90^\circ$

