



Pearson
Edexcel

Examiners' Report
Principal Examiner Feedback

Summer 2023

Pearson Edexcel GCE
In Mathematics (9MA0)
Paper 02 Pure Mathematics

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2023

Publications Code 9MA0_02_2306_ER*

All the material in this publication is copyright

© Pearson Education Ltd 2023

General

This paper offered plenty of opportunity for students to show what they had learnt. The early questions were accessible to candidates of all abilities and this was reflected by candidates regularly scoring full marks on several of the early questions. The longer, later questions provided suitable challenge for stronger candidates but also gave opportunities for restarts for students who struggled with earlier parts of questions.

There were particular instances where marks were lost unnecessarily and the following observations should be helpful to students:

- answers are sometimes not given in the form required, e.g. straight line equations such as in questions 4(b) and 9(b)
- values are sometimes not given as exact or to the required accuracy as demanded in the question, e.g. the exact value of R and the value of α to 3 decimal places in 8(a)
- answers not written as printed on the question paper e.g. the “= 0” missing from 3(a)
- answers not given as requested, e.g. the complete equation not given in 4(b) and the partial fractions not written down in 10(a)

These points aside, there were many succinct and elegant solutions to the more demanding problems towards the end of the paper.

Question 1

This question provided an accessible start to the paper for the vast majority of candidates, most of whom picked up at least some of the marks available.

In part (a) most candidates were able to differentiate $f(x)$ twice and were able to earn both marks. If candidates did lose marks it was mainly due to arithmetic or processing errors such as differentiating $3x^2$ to $5x$. The most common error was a slip in the final coefficient of x .

In (b) part (i), most candidates were able to make a successful attempt to solve $f''(x) = 0$ for x . The mark in this part was sometimes lost due to sign errors when rearranging. It was

surprisingly common to see $6x + 4 = 0$ leading to $x = \frac{2}{3}$. A significant number of candidates did

not simplify their answer, but this was condoned as were values which followed correctly from an incorrect second derivative of the correct form.

Part (ii) proved to more of a challenge and many responses highlighted a lack of understanding or awareness of the condition for a function to be concave. A number of blank responses were seen here. Many who attempted this part of the question deduced, or guessed, that it was necessary to consider the sign of the second derivative but were often confused about which sign was required. Others attempted incorrect calculations setting $f'(x) = 0$ and solving the resulting quadratic in x . Occasionally, candidates attempted sketches of $f(x)$ and/or $f'(x)$ in order to determine the range of $f(x)$ which was not required. Others attempted to demonstrate a sign change in attempt to prove that the solution from part (i) was an inflection point. For those employing the correct method, both strict and non-strict inequalities were accepted.

Question 2

For the most part candidates answered this question well. This was dependent on whether they correctly understood the iteration formula.

In part (a) (i), although many scored this first mark, there was a common misunderstanding of how a recurrence relationship works with confusion as to the meaning of n . Rather than term position, it was often confused with u_n so that the incorrect calculation

$u_2 = 35 + 7 \cos\left(\frac{35\pi}{2}\right) - 5(-1)^{35}$ was often seen, giving the correct answer but obviously not

scoring the B mark. Also not uncommonly seen was n replaced by u_{n+1} , with candidates stating

$$u_2 = 40 + 7 \cos\left(\frac{40\pi}{2}\right) - 5(-1)^{40}.$$

The errors stated above also affected the performance of some candidates in part (ii).

Furthermore, a minority of candidates had difficulty with accuracy in the sign of the final term, struggling with the odd/even power of -1 . For example, when calculating u_3 ,

$$u_3 = 40 + 7 \cos\left(\frac{2\pi}{2}\right) - 5(-1)^2 = 40 - 7 + 5 = 38, \text{ was not uncommon. Occasionally,}$$

candidates were seen to be working with degrees rather than radians, also leading to loss of the accuracy mark.

In part (b) (i) most candidates understood the order 4, periodic nature of the recurrence relationship and that $u_5 = u_1 = u_{4k-3}$, $k \in \mathbb{Z}^+$. Occasionally, candidates unnecessarily calculated u_5 , using u_4 in the recurrence relationship, which sometimes led to the mark being lost either if u_4 had an incorrect value or if an error was made in the calculation.

In part (ii) many candidates recognised how to construct the arithmetic calculation to find the required sum. The most common of these attempts were:

$$6(u_1 + u_2 + u_3 + u_4) + u_1 \quad \text{and} \quad 6(u_2 + u_3 + u_4) + 7u_1$$

Credit was given to candidates who used such ideas with incorrect values from earlier in the question for u_3 and u_4 , although those who had a wrong answer in (a)(i) who used their incorrect value for u_2 (rather than the given $u_2 = 40$), lost both marks. It is important that centres make candidates aware that they must use the 'show that' answer in subsequent calculations no matter what they achieve in their working. Occasionally, candidates used

$$\sum_{r=1}^{25} u_r = 6.25(u_1 + u_2 + u_3 + u_4) \text{ but these usually did not add 1 to give a fully correct}$$

method. Some also incorrectly tried summing the series as though the terms formed an arithmetic progression. Some candidates incorrectly multiplied $(u_1 + u_2 + u_3 + u_4)$ by 4 instead of by 6. A minority forgot to add 35 to their total. A very small minority of candidates generated all the terms of the whole sequence and added them together.

Question 3

There were very few non-attempts or zero scores here and this question proved to be a good discriminator with a spread of marks achieved by candidates. Most students managed to score at least one mark out of three in part (a) by correctly applying a single log law, usually the product, division or power rules for logs. Dealing with the constant, 2, proved to be more challenging for many. Those who wrote 2 as $\log_2(4)$ and then applied the product rule for logs on the right-hand-side had most success. Others grouped the three log terms and applied the exponential rule to eliminate the logs. Sometimes there was a lot of working which wasn't always set out as clearly as it could have been. A significant proportion of candidates did succeed, but it was not uncommon to see correct use of a log law followed by incorrect log work in an attempt to fudge the printed result. Some poor responses involved an initial 'cancelling' of logs before any attempt to combine log terms or to raise both sides to the base 2 correctly. There was a minority of candidates who perhaps knew how to proceed but did not show sufficient working to gain full marks in a 'show that' question. Candidates should be reminded that all steps should be shown in this type of question. It was a shame when fully correct log work was followed by an incomplete 'equation' at the end of this part, as missing the '= 0' meant the loss of the final mark here. A small number of candidates who were unsure how to start part (a) used the printed answer and tried to "work backwards" but often made errors leading to incorrect solutions.

In (b) part (i), almost all candidates were able to correctly state the values '6' and ' $-\frac{5}{3}$ ', it was clear that in many cases candidates were making use of their calculators to solve the quadratic equation which was acceptable. Part (ii) however, was answered less reliably. Although most candidates had some understanding of the domain of a log function and knew that $x = -\frac{5}{3}$ was not a valid solution, many were unable to provide reasoning that was precise enough for the mark here. It was common to see, for example, "a log can't be negative", which was not considered sufficiently clear. Others stated simply 'because it is negative' which was also insufficient. A range of responses which engaged specifically with the domain of a log function needing to be positive were considered acceptable such as; 'log of a negative value is undefined/impossible/gives a math error'.

Question 4

Those scoring full marks in this question were in a significant minority of candidates.

In part (a) the majority of candidates used the given information to find the value of A . Those that did not achieve this mark typically gave 85 as an answer. A few students failed to recognise the standard technique of substituting $t = 0$ into the expression for H to find the value of A , often using $t = 1$. Sometimes e^0 was evaluated as e or 0 rather than 1.

Part (b) had mixed outcomes for candidates. Interpretation of the given initial rate of cooling was problematic for many candidates. There was a general lack of appreciation that the *rate of cooling* was a continuous variable which represented $-\frac{dH}{dt}$. Those candidates not recognising

this and using a difference method (similar to those introduced at GCSE) between $t = 0$ and $t = 1$ were restricted to the initial B mark in part (a). The most common attempt at a response for these candidates was setting $H(1) = 85 - 7.5$ and working from there. Of those who recognised the need to first find $\frac{dH}{dt}$ and then substitute $t = 0$, many incorrectly used $\frac{dH}{dt} = 7.5$ rather than

the correct $\frac{dH}{dt} = -7.5$ to express a negative increase in temperature. This led to the loss of the

last mark. Another error was to forget to differentiate the + 30 to zero leaving a constant in their $\frac{dH}{dt}$, only scoring at most the second method mark for substituting $t = 0$ and their A into their

$\frac{dH}{dt}$. Some differentiated incorrectly to obtain $-55Bte^{-Bt}$. Several students missed out on the

final mark despite correct work by not writing out the final equation. It is important that centres make candidates aware that when a question asks for the complete equation for a model that it is given as such and not just the values of any required constants.

Question 5

Part (a) of this question was accessible for the majority of candidates. Most understood that the derivative at a stationary point is zero and were able to achieve both the method mark and the accuracy mark, substituting $x = 3$ into the given derivative and setting it equal to zero.

Occasionally the accuracy mark was lost due to insufficient working following substitution e.g. $2(3)^3 - 9(3)^2 + 5(3) + k = 0$ followed by $k = 12$ thus omitting a required intermediate step. In a 'show that' question candidates need to ensure that they have completely justified the given answer. Occasionally students showed the substitution of $x = 3$ into the derivative, processed the powers and simplified to obtain $k - 12$ and then jumped to $k = 12$. These responses lacked the appreciation that the correct answer had been achieved because the gradient equals 0 at $x = 3$.

There were some rare instances of candidates incorrectly substituting $x = 3$ and $\frac{dy}{dx} = -10$ to

find the value of k . Some candidates automatically integrated the given derivative but then realised that the differentiated expression was required in order to answer part (a) which they then successfully did. Other candidates integrated the expression and then substituted in the value for $k = 12$ to achieve an answer for part (b). Occasionally they then used the answer to part (b) to find $k = 12$, so were unable to access the marks available for this part of the question. A rarely seen alternative was to use algebraic division to divide the given polynomial by the linear factor $(x - 3)$ to achieve a remainder of $-12 + k$, and then equate that remainder to zero. This approach was met with varying degrees of success and students were often unable to complete the algebraic division successfully.

In part (b), candidates generally understood that they were required to integrate the expression and the straightforward integration meant the first method mark was commonly achieved. Most realised that they then needed to find the value of their constant of integration using $x = 3$ and $y = -10$ to determine the point where the curve crossed the y -axis and achieved the second method mark. The majority of students were successful in achieving full marks in this part, with most stating their answer in coordinate form $(0, -28)$ though some candidates left their answer as $c = -28$, not making the link with the y intercept explicit. Very few candidates made numerical slips in their processing to find the value of their constant of integration. Where mistakes were made, a common reason for not scoring the second method mark was due to substituting in a value of 0 instead of -10 for y in the integrated expression. Where the constant of integration had been omitted candidates went directly to substituting $x = 0$. Less commonly, some candidates differentiated rather than integrated and tried to find the roots of the second derivative rather than the y intercept.

Question 6

This question was relatively well attempted. Most candidates attempted both parts and scored highly overall.

In part (a) the majority of candidates correctly identified $10\mathbf{i} + 24\mathbf{j}$, and $50\mathbf{i} + 120\mathbf{j}$. Those who didn't, often mistakenly added the position vectors instead of subtracting them. Most then went on to show that one was a multiple of the other (or that they were both different multiples of the same vector). However, some wrote the vectors the wrong way round for their multiple, which lost the second mark.

Some candidates used other successful methods involving ratios or "gradients".

A common error here was candidates attempting to conclude the vectors were parallel by calculating and using their magnitudes. The most common reason for the loss of a mark in part (a) was with candidates correctly showing that the two vectors were multiples of each other but then not drawing a conclusion afterwards. It is important that candidates state what is shown by their working and provide a conclusion.

In part (b) most candidates correctly identified the lengths of AB , BC , CD and AD , but many left out their calculations by not showing the application of Pythagoras' Theorem. This is an important step that should be included to fully demonstrate where their answers came from.

There was a misunderstanding by some students who thought they could add the 4 vectors of the sides together to calculate a total vector and work out the length of the resulting vector. A few instead found the lengths of the position vectors and added them. Some also assumed that there were two pairs of equal length lines and/or two pairs of parallel lines and just used the two lengths from part (a) then doubled them. Despite this, candidates often achieved the first two marks. The calculation of the average speed was also done well by many candidates. Although several candidates based their calculation on a single lap and arrived at half the correct answer, the majority completed it correctly and clearly showed division by a fraction equivalent to $\frac{5}{60}$

(most commonly $\frac{1}{12}$) or multiplication by its reciprocal. Candidates should be aware of the

general instruction in the rubric to give non-exact answers to 3 significant figures; it was common to see 7 or 7.0 instead of the correct final answer.

Question 7

Almost all candidates attempted to use implicit differentiation with good levels of success. It was somewhat surprising that a significant number of candidates were let down by failing to differentiate the constant term correctly despite, in many cases, managing to correctly carry out some of the more complex steps in the implicit differentiation. It was not uncommon though to

see errors in either the product rule for $2xy$ (often obtaining $\frac{d}{dx}(2xy) = 2\frac{dy}{dx}$ or

$\frac{d}{dx}(2xy) = 2x\frac{dy}{dx}$ or errors in the application of the chain rule when differentiating $3y^2$. A

surprising number of candidates set their first line of working equal to $\frac{dy}{dx}$ and fortunately this

was often ignored in later working and so was condoned. Some candidates however, attempted to incorporate the extra $\frac{dy}{dx}$ term into their manipulations to find an expression for $\frac{dy}{dx}$ and this

was costly. A few candidates struggled to deal with the negative terms in their numerator (or denominator) which led to sign errors in part (b). Others made slips in the rearrangement which lost the final mark. In some cases, there were issues with clarity of notation for the final

statement of $\frac{dy}{dx}$ in terms of the positioning of the minus sign and candidates should be advised

to take care when drawing the vinculum, particularly when the starting term of the numerator is negative to make sure the sign is in the correct place.

In part (b), the majority of candidates were able to use the correct approach to find the equation of the normal and candidates were often able to earn at least some of the marks here even if they had been unsuccessful in part (a). The least successful approach for calculation of the gradient

of the normal tended to involve an attempt to rearrange the expression for $\frac{dy}{dx}$ into $-\frac{dx}{dy}$ prior to

substitution and this seem to be more error-prone than the alternative of substituting in values as a first. Some candidates did not spot that point P on the curve had been given in the question and so didn't use this point in order to find the equation of the line. In some cases, candidates found the equation of the tangent rather than the normal whilst others made arithmetical slips and lost the accuracy mark. Occasionally candidates did not state their equation in the required form which was a shame when it followed correct work.

Question 8

Part (a) of this question was generally answered very well with candidates regularly scoring full marks. Candidates seemed to be well practiced at converting to harmonic form and working in radians. The most successful candidates wrote out the full expansion of $R \cos(\theta - \alpha)$ rather than trying to take a short cut. The common issues included giving a decimal solution for R or rounding the value of α to a lower degree of accuracy than was required in the question. There were a small number of candidates who worked with $\tan \alpha = \frac{2}{8}$ and so lost the method mark and the accuracy mark.

Candidates found part (b) more challenging, with many failing to link areas of mathematics, in this case trigonometry and series. Even where a candidate successfully found the sum,

$S = 9 \cos x + 36 \sin x$, many did not relate it to part (a). Those who did recognise the form were usually able to get $\pm 4.5 \times "R"$, usually $\pm 4.5 \times 2\sqrt{17}$ for part (i). Some seemed to misread the question and stated the minimum value, so lost the accuracy mark. A significant number of candidates found a correct expression in harmonic form but lost both the M1 and A1 as they kept a correct $9\sqrt{17}$ embedded in their expression.

Candidates were slightly more successful in part (ii). Most candidates who stated a correct value for the maximum, also got the value of x correct, more often than not giving it to the same accuracy as in part (a). A few benefitted from this being a follow through mark. Some candidates tried to find a subsequent value for α and were often out by π or 2π , or attempted to rearrange $x - 1.326 = \pi$. A few responses were seen where S was differentiated and set to zero to find a maximum. These candidates often lost the accuracy mark as they found an answer of 37.1. A lot of incorrect responses gave a maximum of 36 when $\sin x = 1$ or 45 from finding $36 + 9$.

Question 9

Most candidates found part (a) challenging but it was usually attempted. There were not many blank responses. Of the four methods listed on the mark scheme, one common successful method was to substitute t in terms of x into $y = 6\ln(t + 3)$ via “completing the square” (way 1). When substituting into $y = 6\ln(t + 3)$, if a correct t equation was created, then this usually led to full marks. Another common approach was to rearrange t in terms of y , usually

successfully as $t = e^{\frac{y}{6}} - 3$ and substitute into $x = t^2 + 6t - 16$ (way 3). When using this method

some candidates had difficulty squaring $e^{\frac{y}{6}}$ meaning they were unable to earn full marks. Some candidates incorrectly set $\sqrt{x + 25} = \sqrt{x} + 5$ thus also losing the accuracy mark. A significant number of candidates factorised $x = t^2 + 6t - 16$ to give $x = (t - 2)(t + 8)$ and then wrongly thought $x = t - 2$. This then often led to $y = 6\ln(x + 5)$ which scored no marks in this part.

Candidates were generally more successful in part (b) than part (a). Many candidates found the y intercept by substituting $t = 2$ into the parametric equation for y rather than substituting $x = 0$ into the Cartesian equation. The first method mark was more challenging. Some candidates

attempted the derivative from the Cartesian equation and they usually reached either $\frac{3}{(x + A)}$ or

$\frac{3}{x}$ whilst others attempted to use the chain rule approach. Some candidates obtained the correct

derivative but failed to substitute $x = 0$. If a candidate achieved the first method mark, they usually went on to achieve the dependent method mark for attempting to find the equation of the tangent. There were very few cases where the candidate used the negative reciprocal of their gradient. The most successful way candidates achieved full marks was differentiating their Cartesian equation. Those that differentiated parametrically were often caught out by arithmetic and algebraic errors or substituted $t = 0$ instead of $t = 2$ and failed to achieve the gradient correctly. For the final answer, some candidates had failed to realise the form of equation the question was asking for so left their answer as “ $y =$ ” or they didn’t have a , b and c as integers. Many candidates who had incorrectly answered (a), were able to go on to achieve full marks in (b) by using parametric differentiation to work out the gradient. There were very few blank responses for this part of the question.

Question 10

Part (a) was generally done well with most candidates not deterred by the occurrence of the “ k ” in the numerator. The majority of students correctly formed the partial fractions and set up the identity in terms of A and B . Once there, most of them correctly went forward by substituting x as 2 and -4 and finding A and B in terms of k . Some set up their partial fractions in the form

$A + \frac{B}{x+4} + \frac{C}{x-2}$ but very rarely managed to achieve $A = 0$. The method of comparing

coefficients and solving two equations in A and B simultaneously was seen very occasionally. Errors were made where candidates had assigned the wrong values for A and B in their final partial fractions. A common sign error was made when substituting in $x = -4$, leading to solving $-12k - 18 = -6A$ to obtain $A = 2k - 3$, instead of $A = 2k + 3$. Some did not fully simplify their expressions for A or B , leaving them as fractions. Another error at this stage generally involved not substituting for x in the $3kx$ term, leaving A and B in terms of x . Some students transposed the $x - 2$ and $x + 4$ at the beginning so lost the B mark but generally followed through correctly and scored the method mark. In a small number of cases, the last mark was lost after candidates correctly found the numerators in terms of k , but did not write down the correct partial fractions.

Very few candidates managed full marks in part (b), although the majority did gain the first method mark for integrating to obtain $\dots \ln(x+4)$ or $\dots \ln(x-2)$. Those who did not recognise the logarithmic form for the integration were unable to score any marks in this part. Some achieved the correct form after first using a formal substitution. Of those who integrated their partial fractions successfully, the vast majority were far from strict in their use of the modulus symbol and many lost marks due to lack of appreciation of its importance. Use of modulus notation for integrating reciprocal functions should be picked up by centres as a teaching point with future cohorts. Students who had written the incorrect term $-(k-3)\ln(-5)$ or even the correct term $-(k-3)\ln|-5|$ after applying the limit $x = -3$, went on to indicate that they believed this term was zero or could just be ignored. Another example of poor practice was missing brackets around the coefficients e.g. $2k + 3\ln(x+4) + k - 3\ln(x-2)$ being surprisingly common. The “invisible” brackets were sometimes recovered but in many cases were not. There were also examples where the brackets around the $(x+4)$ and the $(x-2)$ were also missing. Those with a systematic well-laid out approach for the substitution of limits in both terms and identifying clear subtraction of the lower limit, went on to score 3 or 4 marks in part (b) and commonly full marks for the question. Many candidates however were hampered by their own layout making the correct interpretation of $\ln(-5)$ as $\ln|5|$ even more challenging. The perhaps unfamiliar, exact form needed for k also proved a challenge for even the best candidates. There were some elaborate attempts to achieve an expression for k , some using exponentials, but many gave up along the way. There were, however some very elegant and correct solutions to this question and some different forms of the exact equivalent answer were seen, for example $21\log_5 e - 6$ but most correct answers were given as $\frac{21}{\ln 5} - 6$.

Question 11

Many candidates found this question challenging and there were a significant number of blank responses.

Candidates found part (a) particularly challenging, with many attaining either no marks or the first mark only. The concept of connected rates of change is one that candidates frequently struggle with. A good number found an expression for V and successfully differentiated to get $\frac{dV}{dh} = 200$, although some confused the volume of water with the volume of the tank. Many

candidates failed to recognise that $\frac{dV}{dt} \propto \frac{1}{\sqrt{h}}$ and hence $\frac{dV}{dt} = \frac{k}{\sqrt{h}}$. Of those who had some

concept of the rate being $\frac{dV}{dt}$, many misunderstood the idea of inverse proportion and others

missed the constant of proportionality. Even where candidates had correct expressions for both $\frac{dV}{dh}$ and $\frac{dV}{dt}$, they often failed to use the chain rule correctly to find $\frac{dh}{dt}$. Many candidates

who did link their expressions correctly, achieving $\frac{dh}{dt} = \frac{1}{200} \times \frac{k}{\sqrt{h}}$

or equivalent, stated that this could be expressed as $\frac{\lambda}{\sqrt{h}}$, with $\lambda = \frac{k}{200}$ and so gained all

three marks. Others made no reference to λ and so lost the accuracy mark. A less common, but

correct, approach used $\frac{dV}{dt} = \frac{d(200h)}{dt}$

Approaches to part (b) of the question were very mixed. Many candidates failed to realise that, as the question said, 'use the model to find an equation', they were expected to, for example, separate the variables and use calculus to solve the differential equation. As they were asked to give their answer in a specific form, a large number started from this and found the constants A and B using the given values of t and h . Fortunately, they were able to gain two marks from the special case. Candidates who used calculus rarely had problems integrating \sqrt{h} , although

errors were seen later when substituting into $h^{\frac{3}{2}}$ or in dealing with the $\frac{2}{3}$ when rearranging the

equation. Candidates who started this part with the λ were generally more successful. Some candidates correctly found c and λ , and then rearranged. Others rearranged before making substitutions, stating $\frac{3}{2}\lambda = A$ and $\frac{3}{2}c = B$. A minority lost the second accuracy mark as they

did not give the final equation in the correct form. Some candidates who had correctly found $\frac{dh}{dt} = \frac{k}{200} \times \frac{1}{\sqrt{h}}$ solved the differential equation using $\frac{k}{200}$ in place of λ , in most cases

successfully, although the algebraic manipulation was more tricky. Some candidates who had

made an incorrect attempt at (a) continued with their answer, usually $\frac{dh}{dt} = \frac{1}{200} \times \frac{1}{\sqrt{h}}$, in this

part. As they only had one constant, the 'c', in their expression, they could only attain the first two method marks. It was rare to see the constant of integration missed and basic arithmetical

errors when finding either λ and c or A and B were also relatively infrequent. The alternative approach, rearranging to give $\frac{dt}{dh} = \frac{\sqrt{h}}{\lambda}$ or $\frac{200\sqrt{h}}{k}$ was seen on a few occasions.

The majority of candidates who had a correct form of the equation in part (b) made a good attempt at part (c) and were generally able to gain the method mark. Relatively few lost the accuracy mark due to missing the units in their final answer. A number of candidates used an incorrect value for h , most commonly 1000, and so lost both marks.

Question 12

In part (a), most candidates were successful in understanding the requirement to substitute zero into N_A and N_B . There were some processing errors resulting in incorrect values for N_A and N_B . A significant number failed to score the method mark by not subtracting their values. There were a few candidates that did not recognise the question was scaled in thousands and left a final answer of 5.

In part (b), a large majority of candidates were able to deduce that $T = 3$ and most explanations were around the idea of the values increasing after this point. The next most common explanation centred on the graph being a minimum or having “turning point” at this location. There were also candidates who stated the gradient has increased or was positive.

In part (c), a significant number of candidates failed to identify either of the correct equations. Many were able to identify and solve correctly at least one equation to find one critical value for t . A common error often followed their correct equation of $5 = 3t$ when they wrote the value of t as $3/5$ and similarly, but less common, $13 = 3t$ leading to $t = \frac{3}{13}$. Some students kept the modulus

signs within their equations and ignored them at various stages in their working which often led to incorrect equations. There was an array of methods seen trying to negotiate the modulus signs, and $|A| + |B| = |A + B|$ was seen a few times and $|A| + |B| = k \Rightarrow |A|^2 + |B|^2 = k^2$ was also seen.

Those who found two values for t were generally able to score the 2nd follow through accuracy mark for choosing the outside region for their critical values. The biggest challenge was the final accuracy mark as quite a few candidates did not write the inequalities using set notation. Those who attempted set notation usually were unable to write their answer correctly

in that form, some confusing \cup with \cap . A common answer was $\left\{ t < \frac{5}{3} \cup t > \frac{13}{3} \right\}$ which

was condoned for this final mark.

Those who attempted the squaring method in this part had minimal success depending on when they chose to square.

Many candidates were successful in part (d) even if they had not progressed with the other parts of the question. The majority of candidates correctly referenced the subscribers will become negative to obtain this B mark. Where a candidate attempted this part but did not obtain this mark, it was usually due to not making a reference to the number of subscribers becoming negative, saying it goes to zero or that it is linear or decreasing. Several candidates thought it couldn't go above 8000 subscribers.

Question 13

Most candidates attempted at least part (a) of this question, but some found the work more demanding in (b) and (c) and made little progress in either part.

Despite this, there were many fully correct or nearly fully correct attempts at the whole question.

In part (a), most candidates successfully factorised out 3^{-2} and attempted the correct resulting expansion of $\left(1 + \frac{x}{3}\right)^{-2}$. There were some arithmetic slips, but generally candidates often reached the correct answer. A common error, however, was failing to factorise properly, reaching $3^{-2}(1+x)^{-2}$. There were a number of attempts at direct expansion. Some of these used non-standard binomial coefficients involving negative integers. To achieve credit for using these, candidates needed to demonstrate the meaning of their non-standard notation. Very occasionally candidates multiplied their correct simplified expansion by 9 or 27 at the end. This still received full marks as examiners could ignore subsequent working following seeing a correct simplified answer.

In part (b), many candidates correctly multiplied their expansion by $6x$ and then integrated using the limits of 0.2 and 0.4 reaching 0.03304, gaining full marks. A common error, however, was not giving the final answer to the required accuracy of four significant figures, thus losing the final accuracy mark. Those candidates who had expanded part (a) to more than three terms also lost this final accuracy mark as it led to a different approximation. A more complicated and unnecessary method involving integration by parts was used by a small minority of candidates. Candidates also often tried inappropriate approaches like dividing $6x$ by their expansion followed by a logarithmic integral. A minority of candidates, despite the warning at the start of the question, clearly used calculator technology to obtain 0.032865, rather than the answer of 0.03304 required. Others made inappropriate attempts at the trapezium rule, thus gaining no credit as algebraic integration was required.

In part (c), the commonest methods of integration adopted were substitution, using either $u = x + 3$ or $u = (x + 3)^2$, integration by parts, integration after applying partial fractions and integration using the reverse chain rule after writing the integrand in appropriate form. Candidates had varying success with all of these approaches, gaining a variety of marks. Integration by parts seemed to lead to most arithmetical mistakes. Errors resulted from using parts ‘the wrong way round’ and others integrated the $(3 + x)^{-2}$ term incorrectly. When

integrating by substitution some students incorrectly thought that $-\frac{18}{u^2}$ would integrate to a

natural logarithm. There was some confusion with the limits; some candidates did not change the limits to “ u ” values and other who did evaluate new limits for their substitution switched back to an expression in x before substituting their changed limits. When using partial fractions some students struggled with the initial format for the partial fractions and it was not uncommon

to see partial fractions of the form $\frac{A}{3+x} + \frac{B}{3+x}$. There was a number of candidates who

either did not attempt this part of the question at all or who lacked a correct strategy. There were responses where students attempted to integrate numerator and denominator separately, e.g. $6x$

to $3x^2$ and $\frac{1}{(3+x)^2}$ to a variation of $\ln(3+x)^2$, and proceeded to multiply both results. A

number of candidates were awarded just the first mark as their overall problem solving strategy was correct despite the fact they were unable to fully implement it. There were also many candidates who could carry out the work in this part completely successfully.

Question 14

Part (a) was at least started by many but when students failed to make any progress towards the result, they tended not to attempt part (b).

In part (a) the vast majority of candidates scored the first mark for using a correct trigonometric identity. The M and A marks were often both given or both lost, as candidates explored multiple manipulative steps to reach the final form for the equation. In many cases this involved expanding the brackets, using a substitution for $\sin 2\theta$, but later putting back into $\sin 2\theta$ form i.e., undoing the manipulation that had been introduced. Whilst in many cases this resulted in the candidate getting there in the end, there were many examples of multiple attempts or unnecessarily long solutions. The most efficient methods did not involve multiplying out the bracket on the LHS which seemed to be the default first step for most students. A common mistake was to suddenly put the RHS equal to zero so it disappeared rather than being subtracted on the LHS. For those that were successful, there were a variety of approaches as well as variety in the number of steps required to reach the correct final form, with some methods considerably more efficient than others. Arithmetic errors were commonly seen in this question. Many successful candidates attempted to find a factor of $\sin 2\theta$ on both sides of the equation before proceeding to collecting these on the same side. Another typical successful method involved writing all the terms in terms of $\sin \theta$ and $\cos \theta$, collecting all terms on the same side, and then taking a factor of $2\sin \theta \cos \theta$. However, many candidates who expanded tended to forget terms when expanding or factorising, leading to an expression in an incorrect format. Replacing $\tan \theta$ and $\sin 2\theta$ was usually handled well, as was replacing $1 + \tan^2 \theta$ with $\sec^2 \theta$ and then $\frac{1}{\cos^2 \theta}$ but many candidates didn't follow the manipulation to the end and were

unsure what to do with the $\sin 2\theta$ on the right hand side, so either left it on the right hand side and lost both the M and A marks, or moved it to the left but then incorrectly dealt with it when combined with the quadratic formed on that side.

Part (b) was generally done well by the candidates who managed to obtain a quadratic in part (a), with many candidates correctly solving to get 360 or 540, and then solving their quadratic. Those who found a solution to their quadratic also tended to find solutions based on that within the specified set of values. Where marks were lost, it was often by incorrectly including extra values, in particular the value of 450 degrees which had been specifically excluded in the question.

Question 15

A number of candidates lost all marks on this question by not attempting any solution.

Most candidates correctly expanded the given expression and used $\sin^2 x + \cos^2 x = 1$ to simplify and score the first two marks. A small number incorrectly used $(\sin x - \cos x)^2 = \sin^2 x - \cos^2 x$, scoring no marks. Others attempted to substitute values of 90 and 180 into the given expression instead of using algebraic manipulation.

Whilst most candidates scored the first two marks, only the very best were able to score the final mark, with most candidates failing to offer a convincing reason for a contradiction. Candidates who drew a graph of $\sin x$ and $\cos x$ or $\sin 2x$ to support their understanding tended to score well. Successful candidates typically explained that $2x$ must be between 180 and 360, and hence $\sin 2x$ must be negative, which is a contradiction with their inequality. Those that worked with $\sin x \cos x$ were generally more successful in achieving this mark than those who chose to use $\sin 2x$. However, it was clear that not all students understood what an obtuse angle was. Another typical correct conclusion involved candidates stating if x is obtuse, $\sin x$ must be positive and $\cos x$ must be negative, leading to a negative product and a contradiction. Of those who were able to offer a reason for the contradiction, some still lost the final A mark for failing to then reach a conclusion that the original statement was true or because of errors in the proof such as missing x 's or mixed variables.

